

Positive gravitational energy in arbitrary dimensions

Yvonne Choquet-Bruhat

July 22, 2011

Résumé

On démontre un théorème d'énergie gravitationnelle positive en dimension quelconque utilisant seulement des spineurs liés au groupe $Spin(n)$ sur une section d'espace Riemannienne (M^n, g)

Abstract

We present a streamlined, complete proof, valid in arbitrary space dimension n , and using only spinors on the oriented Riemannian space $(M^n; g)$, of the positive energy theorem in General Relativity.

Version française abrégée.

Un espace-temps Einsteinien est une variété Lorentzienne $(\mathbf{M}^{n+1}, \mathbf{g})$ qui satisfait les équations d'Einstein

$$\mathbf{R}_{\alpha\beta} - \frac{1}{2}\mathbf{g}_{\alpha\beta}\mathbf{R} = T_{\alpha\beta};$$

on supposera qu'il satisfait la condition d'énergie dominante, $u_\alpha T^{\alpha\beta}$ temporel pour tout vecteur temporel u . Sur chaque section spatiale M^n la métrique induite g et la courbure extrinsèque K satisfont les contraintes qui s'écrivent dans un repère orthonormé d'axes e_i tangents et e_0 orthogonal à M^n

$$\mathbf{R}_{0j} \equiv \partial_j K_h^h - D_h K^h_j = T_{0j} \quad (0.1)$$

$$\mathbf{S}_{00} \equiv \frac{1}{2}\{R - |K|^2 + (\text{tr} K)^2\} = T_{00}. \quad (0.2)$$

On suppose que M^n est l'union d'un compact W et d'un nombre fini N d'ensembles Ω_I , appelés bouts (ends), diffeomorphes au complément d'une

boule de R^n . On utilise une partition lisse f_I, f_K de l'unité sur M^n de supports contenus dans un Ω_I ou un ouvert W_K difféomorphe à une boule de R^n , l'union (finie) des W_K recouvrant W . Tous ces ouverts sont munis de coordonnées locales x^i et de la métrique Euclidienne $e \equiv \eta_{ij}dx^i dx^j$. Un tenseur u sur M^n est une somme de tenseurs $u_I = f_I u$, $u_K = f_K u$. On utilise la métrique euclidienne pour définir les normes de ces tenseurs. Un espace de Banach C_β^k ou de Hilbert $H_{s,\delta}$ d'un tenseur u sur M^n est défini à l'aide du *sup* ou de la somme des normes des tenseurs u_I et u_K , des choix différents de partition de l'unité donnent des normes équivalentes. La variété Riemannienne (M^n, g) est dite asymptotiquement Euclidienne (A.E) si

$$h := g - \underline{g} \in H_{s,\delta} \cap C_{n-2}^1, \quad s > \frac{n}{2} + 1, \quad \frac{n}{2} - 2 > \delta > -\frac{n}{2}, \quad (0.3)$$

où $\underline{g}$ est une métrique lisse identique dans chaque Ω_I à la métrique Euclidienne e .

La définition de masses gravitationnelles m_I et de moments p_I , dits ADM, d'un espace-temps with A.E. section (M^n, g) and $K \in H_{s-1,\delta+1}$ provient de la formulation Hamiltonienne des équations d'Einstein, on a dans Ω_I

$$m_I := \lim_{r \rightarrow \infty} \frac{1}{2} \int_{S_r^{n-1}} \left(\frac{\partial g_{ij}}{\partial x^j} - \frac{\partial g_{jj}}{\partial x^i} \right) n_i \mu_e, \quad r \equiv \left\{ \sum_i (x^i)^2 \right\}^{\frac{1}{2}}, \quad (0.4)$$

$$p_I^h := \lim_{r \rightarrow \infty} \int_{S_r^{n-1}} P^{ih} n_i \mu_e, \quad P^{ih} := K^{ih} - g^{ih} \text{tr} K. \quad (0.5)$$

On reprend l'idée spinorielle de Witten pour démontrer, mais en utilisant seulement un spineur sur M^n lié à l'algèbre de Clifford $\mathcal{Cl}(n)$, la positivité de la masse d'un espace temps Einsteinien quand $R \geq 0$, donc sous la condition d'énergie dominante, quand M^n est une hypersurface maximale. Une formulation simple liée au moment P , qui ne fait intervenir que la même sorte de spineurs sur M^n , permet de montrer que $m_I \geq |p_I|$ donc $m \geq |p|$ sans condition sur R ni autre hypothèse sur les sources. Les démonstrations reposent sur un théorème d'existence pour la solution d'une équation de Dirac complétée, elliptique, sur une variété asymptotiquement Euclidienne.

English version

1 Introduction

The most elegant and convincing proof of the positive energy theorem is by using spinors, as did Witten¹ in the case $n = 3$ inspired by heuristic works of Deser and Grisaru originating from supergravity. The aim of this Note is to present a streamlined, complete proof, valid in arbitrary space dimension n , and using only spinors on the oriented Riemannian space $(M^n; g)$, without invoking spacetime spinors.

We first give the notations and the definitions we use.

2 Definitions.

2.1 Asymptotically Euclidean space.

M^n is a smooth manifold union of a compact set W and a finite number of sets Ω_I , diffeomorphic to the complement of a ball in R^n . One covers W by a finite number of open sets W_K each diffeomorphic to a ball in R^n . We denote by x^i local coordinates for a domain Ω_I or W_K . We set $r := \sum (x^i)^2\}^{\frac{1}{2}}$ and take $r_0 > 0$ such that $\Omega_I := \{r > r_0\}$, $\Omega_I \cap W_K = \emptyset$ if $r < 2r_0$. We consider a preparation of M^n , i.e a smooth partition of unity, f_I, f_K , with support in W_k , f_I support in Ω_I and $f_I = 1$ for $r > 2r_0$. The Riemannian metric g is continuous and uniformly bounded above and below in each Ω_I , W_K by constant positive definite quadratic forms. A tensor field u on M^n is written as $u \equiv \sum_I u_I + \sum_K u_K$ with $u_I := f_I u$, $u_K := f_K u$. Norms on spaces of tensor fields are defined through their components in the Ω_I , W_K , each endowed with the Euclidean metric $e := \eta_{ij} dx^i dx^j \equiv \sum (dx^i)^2$, with pointwise norm $|\cdot|$ and volume element μ_e . We use the Banach and Hilbert spaces C_β^k and $H_{s,\delta}$ with norms

$$\|u\|_{C_\beta^k} \equiv \sup_{I,K} \{ \sup_{\Omega_I} (r^{\beta+k} |\underline{D}^k u_I|), \sup_{W_K} |\underline{D}^k u_K| \}, \quad \underline{D}^k := \frac{\partial^k}{\partial x^{i_1} \dots \partial x^{i_k}}. \quad (2.1)$$

$$\|u\|_{H_{s,\delta}}^2 := \sum_{I=1,\dots,N} \int_{\Omega_I} \sum_{0 \leq k \leq s} r^{2(k+\delta)} |\underline{D}^k u_I|^2 \mu_e + \sum_{K=1,\dots,N'} \int_{W_K} \sum_{0 \leq k \leq s} |\underline{D}^k u_K|^2 \mu_e \quad (2.2)$$

¹For references prior to 1983 one can consult my survey on positive energy theorems for les Houches 1983 school reproduced in Y.Choquet-Bruhat 'General Relativity and the Einstein equations', Oxford University press 2009.

Different preparations of M^n give equivalent norms. A Riemannian manifold (M^n, g) is called asymptotically Euclidean (A.E) if

$$h_I := f_I(g - e) \in H_{s,\delta} \cap C_{n-2}^1, \quad f_K g \in H_s, \quad s > \frac{n}{2} + 1, \quad \frac{n}{2} - 2 > \delta > -\frac{n}{2}. \quad (2.3)$$

It can be proved (using the fact that $H_{s,\delta}$ is an algebra if $s > \frac{n}{2}, \delta > -\frac{n}{2}$) that an A.E (M^n, g) admits in each end Ω_I an orthonormal coframe

$$\theta^j := a_i^j dx^i \quad a_i^j = \delta_i^j + \frac{1}{2} \lambda_i^j, \quad \lambda_i^j \in H_{s,\delta} \cap C_{n-2}^1. \quad (2.4)$$

In the following, components in the coordinates x^i are underlined. In Ω_I it holds that

$$\underline{g}_{ij} \equiv \sum_h a_i^h a_j^h \equiv \eta_{ij} + \underline{h}_{ij}, \quad \eta_{ij} := \delta_i^j, \quad \underline{h}_{ij} \equiv \frac{1}{2}(\lambda_i^j + \lambda_j^i) + \frac{1}{4} \sum_h \lambda_j^h \lambda_i^h, \quad \lambda_j^h \lambda_i^h \in H_{s,2\delta+\frac{n}{2}} \cap C_{2n-4}^1. \quad (2.5)$$

The rotation coefficients c_{ij}^h of the coframe θ^h are, with (b_i^j) the matrix inverse of (a_j^i) and ∂_i the Pfaff derivative with respect to θ^i , $d\theta^h \equiv \frac{1}{2} c_{ij}^h \theta^i \wedge \theta^j$,

$$c_{ij}^h \equiv b_j^k \partial_i \lambda_k^h - b_i^k \partial_j \lambda_k^h \equiv \frac{1}{2}(\partial_i \lambda_j^h - \partial_j \lambda_i^h) + \chi_{ij}^h, \quad \chi_{ij}^h \in H_{s-1,2\delta+1+\frac{n}{2}} \quad (2.6)$$

We choose the coframe such that

$$\partial_i(\lambda_i^j - \lambda_j^i) \in H_{s-1,2\delta+1+\frac{n}{2}}$$

The components $\omega_{i,jh} \equiv \frac{1}{2}(-c_{jh}^i + c_{ij}^h - c_{ih}^j)$ of the Riemannian connection ω in the coframe θ^i are then computed to be

$$\omega_{i,hj} \equiv \frac{1}{2} \underline{\partial}_h \underline{h}_{ij} - \frac{1}{2} \underline{\partial}_j \underline{h}_{ih} + \zeta_{i,hj}, \quad \zeta_{i,hj} \in H_{s-1,2\delta+1+\frac{n}{2}}. \quad (2.7)$$

2.2 Global mass m and linear momentum p

We say that (M^n, g, K) is A.E. if (M^n, g) is A.E. and $K \in H_{s-1,\delta+1} \cap C_{n-1}^0$. The mass m and linear momentum p associated to an end Ω define a space-time vector $\mathbf{E}$ with components

$$E^0 := m := \lim_{r \rightarrow \infty} \frac{1}{2} \int_{S_r^{n-1}} (\underline{\partial}_j \underline{h}_{ij} - \underline{\partial}_i \underline{h}_{jj}) n_i \mu_{\bar{g}}, \quad E^h := p^h := \lim_{r \rightarrow \infty} \int_{S_r^{n-1}} P^{ih} n_i \mu_{\bar{g}}. \quad (2.8)$$

The uniform bound of n_i and the equivalence of $\mu_{\bar{g}}$ with $r^{n-1}\mu_{S_1^{n-1}}$ show that the limits exist.

We always assume that the constraints 0.1 and 0.2 are satisfied on M^n and that T obeys the dominant energy condition.

3 Spinor fields and Dirac operator.

The gamma matrices associated with an orthonormal coframe θ^i of g at $x \in M^n$ are linear endomorphisms of a complex vector space S of dimension $p := 2^{\lfloor n/2 \rfloor}$ which satisfy the identities

$$\gamma_i \gamma_j + \gamma_j \gamma_i = 2\eta_{ij} I_p, \quad i, j = 1, \dots, n, \quad I_p \text{ identity matrix} \quad (3.1)$$

The γ_i are chosen hermitian, i.e. $\gamma_i = \tilde{\gamma}_i$, as is possible for an $O(n)$ group.

The spinor group $Spin(n)$, double covering of $SO(n)$, can be realized by the group of invertible linear maps Λ of S which satisfy, with $O := (O_i^j)$ a $n \times n$ orthogonal matrix

$$\Lambda \gamma^i \Lambda^{-1} = O_j^i \gamma^j \quad \text{and} \quad \det \Lambda = 1. \quad (3.2)$$

In a subset Ω_I or W_K with a given field ρ_0 of orthonormal frames a spinor field ψ is represented by a mapping $(x^i) \mapsto \underline{\psi}(x^i) \in S$. Under an $O \in SO(n)$ change of frame, $\rho = O\rho_0$ the spinor ψ becomes represented by $\underline{\psi}' = \Lambda \underline{\psi}$ where some choice has been made for the correspondence between $\underline{\Lambda}$ and $\underline{O}$. This can be made consistently on M^n if it admits a spin structure; that is, a homomorphism of a $Spin(n)$ principal bundle P_{Spin_n} onto the principal bundle of oriented orthonormal frames. It is a topological property of M^n , the vanishing of its second Stiefel-Whitney class, always true for an orientable M^3 . A spinor field on M^n is then a section of a vector bundle $\Psi_{Spin(n)}$ associated with $P_{Spin(n)}$, with base M^n and typical fiber S . To a space of spinors corresponds a space of cospinors, replacing S by the adjoint (complex dual) vector space $\tilde{S}$ and the change of representation by $\underline{\phi}' = \underline{\phi} \Lambda^{-1}$. Using dual frames e_A of S and θ^A of $\tilde{S}$ we have $\psi \equiv \psi^A e_A$, $\phi = \theta^A \phi_A$, $A = 1, \dots, p$, we denote the duality relation by

$$\phi\psi \equiv (\phi, \psi) := \phi_A \psi^A, \quad \text{a frame independent scalar.}$$

By 3.2 $\tilde{\psi}$ represented by $\tilde{\psi}_A := (\psi^A)^*$ is a cospinor if ψ is a spinor, $|\psi|^2 \equiv \tilde{\psi}\psi$ is positive definite.

A *spin connection* σ on (M^n, g) is deduced from an $O(n)$ connection ω by the isomorphism between the Lie algebras of $O(n)$ and $Spin(n)$ obtained by differentiation of 3.2, it is represented in each domain of the preparation by

$$\sigma_i \equiv \frac{1}{4} \gamma^h \gamma^k \omega_{i,hk}, \quad i = 1, \dots, n. \quad (3.3)$$

The covariant derivative of a spinor ψ , resp. cospinor ϕ , is a covariant vector spinor, resp. cospinor, with components in the frames $\theta^i \otimes e_A$, resp. $\theta^i \otimes \theta^A$,

$$(D_i \psi)^A \equiv \partial_i \psi^A + (\sigma_i \psi)^A, \quad (D_i \phi)_A \equiv \partial_i \phi_A - (\phi \sigma_i)_A.$$

The hermiticity of γ_i and 3.1 show that $\tilde{\sigma}_i = -\sigma_i$, hence $\widetilde{D_i \psi} \equiv D_i \tilde{\psi}$.

The Riemannian connection together with the spin connection define a first order derivation operator mapping tensor-spinor-cospinor fields into tensor- spinor- cospinor fields with one more covariant index. *The gamma matrices are the components of a vector-spinor-cospinor which has covariant derivative zero.*

The spin curvature ρ is a 2-tensor- spinor -cospinor, image by the mapping of Lie algebras of the curvature tensor of g . The Ricci identity for spinors reads

$$D_i D_j \psi - D_j D_i \psi \equiv \rho_{ij} \psi \quad \text{with} \quad \rho_{ij} := \frac{1}{4} R_{ij,hk} \gamma^h \gamma^k. \quad (3.4)$$

The Dirac operator on sections of the vector bundle $\Psi(n)$ reads locally

$$\mathcal{D} \psi \equiv \gamma^i D_i \psi \equiv \gamma^i (\partial_i \psi + \frac{1}{4} \omega_{i,hk} \gamma^h \gamma^k \psi), \quad \text{hence} \quad \widetilde{\mathcal{D} \psi} \equiv D_i \tilde{\psi} \gamma^i. \quad (3.5)$$

The algebraic Bianchi identity together with 3.1 and 3.4 lead to the formula²

$$\mathcal{D}^2 \psi \equiv \eta^{ij} D_i D_j \psi + \frac{1}{2} \gamma^i \gamma^j \rho_{ij} \psi \equiv \eta^{ij} D_i D_j \psi - \frac{1}{4} R \psi. \quad (3.6)$$

The Dirac operator is a first order linear operator with principal symbol $(\eta^{ij} \xi_i \xi_j)^{\frac{p}{2}}$, hence elliptic. Weighted Sobolev spaces for spinor fields on a prepared M^n are defined as for tensor fields after setting $\psi = \sum (f_I \psi + f_K \psi)$ and using representations $\underline{\psi}$. A known theorem³ gives:

²See for instance A. Lichnerowicz Bull. Soc. Math. France 92, 1964 p. 11-100

³Y. Choquet-Bruhat and D. Christodoulou, Acta Mathematica 146, 1981..

Theorem 1 *On an A.E. (M^n, g) the Dirac operator is a Fredholm operator from spinors in $H_{s,\delta}$ to spinors in $H_{s-1,\delta+1}$, it is an isomorphism if injective. The same is true of $\mathcal{D}\psi + f\psi$ if f is a bounded linear map from spinors in $H_{s,\delta}$ to spinors in $H_{s-1,\delta+1}$.*

4 Gravitational mass.

We prove for arbitrary $n > 2$ the fundamental fact used by Witten for $n = 3$.

Theorem 2 *Let (M^n, g) be A.E. The mass m of an end Ω_I is equal to*

$$\lim_{r \rightarrow \infty} \int_{S_r^{n-1}} \mathcal{U}_0^i n_i \mu_{\bar{g}} = \frac{m}{2}, \quad \mathcal{U}_0^i := \operatorname{Re} \{ \tilde{\psi}_0 (\eta^{ij} - \gamma^i \gamma^j) \sigma_j \psi_0 \}, \quad (4.1)$$

with S_r^{n-1} the submanifold of the end Ω_I with equation $\{\sum (x^i)^2\}^{\frac{1}{2}} = r$, n_i its unit normal, $\mu_{\bar{g}}$ the volume element induced by g and ψ_0 a spinor constant in Ω_I (i.e. $\frac{\partial \psi_0}{\partial x^i} = 0$) and $|\psi_0| = 1$.

Proof. We first remark that, using $\gamma^i = \tilde{\gamma}^i$ and $\tilde{\sigma}_i = -\sigma_i$, one finds

$$\operatorname{Re}(\tilde{\psi}_0 \eta^{ij} \sigma_j \psi_0) \equiv \frac{1}{2} \tilde{\psi}_0 \eta^{ij} (\sigma_j + \tilde{\sigma}_j) \psi_0 \equiv 0.$$

The definition 3.3 of σ_j then implies

$$\mathcal{U}_0^i = \frac{1}{8} \sum_{j,h,k} \tilde{\psi}_0 \omega_{j,hk} (\gamma^i \gamma^j \gamma^h \gamma^k - \gamma^h \gamma^k \gamma^j \gamma^i) \psi_0.$$

The property 2.7 of ω on A.E. (M^n, g) shows that $r^{n-1}(\omega_{j,hk} - \frac{1}{2} \underline{\partial}_h h_{jk} + \frac{1}{2} \underline{\partial}_k h_{jh})$ tends uniformly to zero as r tends to infinity. The uniform bound of n_i and the equivalence of $\mu_{\bar{g}}$ with $r^{n-1} \mu_{S_1^{n-1}}$ show that the limit in 4.1 exists. Calculations⁴ using 3.1, 2.7 and the symmetry of $\underline{\partial}_k h_{hj}$ in h and j lead to

$$\lim_{r \rightarrow \infty} \int_{S_r^{n-1}} \mathcal{U}_0^i n_i \mu_{\bar{g}} = \frac{1}{2} \lim_{r \rightarrow \infty} \int_{S_r^{n-1}} \omega_{j,ij} |\psi_0|^2 n_i \mu_{\bar{g}} = \frac{1}{2} m \quad \text{if } |\psi_0|^2 = 1.$$

■

⁴Similar to those done by P. Chrusciel in his Krakow lectures on Energy in General Relativity.

To study the positivity of the mass one defines a vector $\mathcal{U}^i$ on M^n such that the integrals on S_r^{n-1} of $\mathcal{U}^i$ and $\mathcal{U}_0^i$ have the same limit when $r \rightarrow \infty$. The Stokes formula applied to the integral of the divergence of $\mathcal{U}^i$ will give information on this limit. We set

$$\mathcal{U}^i := \mathcal{R}e\{\tilde{\psi}(\eta^{ij}D_j\psi - \gamma^i\gamma^jD_j\psi)\}. \quad (4.2)$$

Lemma 3 *On an A.E manifold (M^n, g) it holds that*

1.

$$D_i\mathcal{U}^i \geq 0 \quad \text{if } R \geq 0 \quad \text{and} \quad \mathcal{D}\psi = 0.$$

2. *If $\psi = \psi_0 + \psi_1$ with $\underline{\partial}_i\psi_0 = 0$ in Ω_I and $\psi_1 \in H_{s,\delta}$ then in Ω_I*

$$\lim_{r \rightarrow \infty} \int_{S_r^{n-1}} \mathcal{U}_0^i n_i \mu_{\bar{g}} = \lim_{r \rightarrow \infty} \int_{S_r^n} \mathcal{U}^i n_i \mu_{\bar{g}}. \quad (4.3)$$

Proof. 1. By elementary computation, using $D_i\tilde{\psi} = \widetilde{D_i\psi}$ and the identity 3.6 one finds

$$D_i\mathcal{U}^i \equiv |D\psi|^2 - |\mathcal{D}\psi|^2 + \frac{1}{4}R|\psi|^2 \quad (4.4)$$

Therefore $D_i\mathcal{U}^i \geq 0$ if $R \geq 0$ and ψ satisfies the equation $\mathcal{D}\psi = 0$.

2. To study the limit of the integral on S_r^{n-1} of $\mathcal{U}^i$ when $\psi = \psi_0 + \psi_1$ we write

$$\mathcal{U}^i = \mathcal{U}_0^i + \frac{1}{2}\mathcal{R}e\{\tilde{\psi}_0[\gamma^j, \gamma^i]D_j\psi_1 + \tilde{\psi}_1[\gamma^j, \gamma^i]D_j\psi)\}. \quad (4.5)$$

Hence $D_i\mathcal{U}^i = D_i\mathcal{U}_0^i + D_i\mathcal{V}^i$,

$$\mathcal{V}^i \equiv \frac{1}{2}\mathcal{R}e\{\tilde{\psi}_0[\gamma^j, \gamma^i]D_j\psi_1 + \tilde{\psi}_1[\gamma^j, \gamma^i]D_j\psi)\} \quad (4.6)$$

Embedding and multiplication properties of Sobolev spaces give

$$\tilde{\psi}_1[\gamma^j, \gamma^i]D_j\psi \in H_{s,\delta} \times \{C_{n-1}^1 \cap H_{s-1,\delta+1}\} \subset H_{s-1,2\delta+1+\frac{n}{2}} \subset C_\beta^0$$

$$\beta < 2\delta + 1 + \frac{n}{2} + \frac{n}{2} < 2n - 3.$$

To estimate the other term one remarks that

$$D_i\{\tilde{\psi}_0[\gamma^j, \gamma^i]D_j\psi_1\} \equiv D_iD_j\{\tilde{\psi}_0[\gamma^j, \gamma^i]\psi_1\} - D_i\{D_j\tilde{\psi}_0[\gamma^j, \gamma^i]\psi_1\} \quad (4.7)$$

the first parenthesis is an antisymmetric 2-tensor hence its double divergence $D_i D_j$ is identically zero. The second parenthesis is

$$D_j \tilde{\psi}_0 [\gamma^j, \gamma^i] \psi_1 \in C_{n-1}^0 \times H_{s,\delta} \subset C_\beta^0.$$

The Stokes formula implies, with $M_r^n := M^n - \{\Omega_I \cap \sum (x^i)^2 \geq r^2\}$

$$\int_{M_r^n} D_i \mathcal{U}^i \mu_g = \int_{S_r^{n-1}} \mathcal{U}^i n_i \mu_{\bar{g}} = \int_{S_r^{n-1}} (\mathcal{U}_0^i + \mathcal{V}^i) n_i \mu_{\bar{g}}. \quad (4.8)$$

The fall off properties found for $\mathcal{V}^i$ complete the proof. ■

Lemma 4 *If (M^n, g) is A.E. $R \geq 0$ and ψ_0 is a smooth spinor constant in Ω_I and zero in the other ends there exists on M^n a spinor $\psi \equiv \psi_0 + \psi_1$, such that $\mathcal{D}\psi = 0$, $\psi_1 \in H_{s,\delta}$.*

Proof. The hypotheses made on ψ_0 show that $\mathcal{D}\psi_0 \in H_{s-1,\delta+1}$. Theorem 1 implies the existence of ψ_1 . ■

The lemmas imply that $m \geq 0$ if $R \geq 0$, that is if (M^n, g) is a maximal submanifold of $(\mathbf{M}^{n+1}, \mathbf{g})$; equivalently, if the pointwise gravitational momentum P on M^n has a vanishing trace. We will now lift this restriction, proving moreover that $m \geq |p|$.

5 Positive energy.

We define a real vector $\mathcal{P}$ on an A.E. (M^n, g, K) by⁵

$$\mathcal{P}^i := \frac{1}{2} \tilde{\psi} \gamma_h P^{ih} \psi \equiv \frac{1}{2} \tilde{\psi} (\gamma_h K^{ih} - \gamma^i \gamma^j \gamma^h K_{jh}) \psi, \quad P^{ih} = K^{ih} - \delta^{ih} \text{tr} K$$

If ψ_0 is as before a smooth spinor constant in one end of M^n and zero in the other ends and ψ is a spinor on M^n such that $\psi - \psi_0 \in C_\beta^0$, $\beta > 0$, then

$$\lim_{r \rightarrow \infty} \int_{S_r^{n-1}} \mathcal{P}^i n_i \mu_{\bar{g}} = \frac{1}{2} \tilde{\psi}_0 \gamma_h p^h \psi_0, \quad p^h := \lim_{r \rightarrow \infty} \int_{S_r^{n-1}} P^{ih} n_i \mu_{\bar{g}}. \quad (5.1)$$

It is elementary to check using the properties of the γ 's that $\gamma_h p^h$ is an hermitian matrix with eigenvalues $\pm |p|$. If we choose for ψ_0 an eigenvector of the eigenvalue $-|p|$ we then have

$$\tilde{\psi}_0 \gamma_h p^h \psi_0 = -|\psi_0|^2 |p|. \quad (5.2)$$

⁵Remark that we do not to introduce a matrix γ_0

To estimate the limit 5.1. we use again the Stokes formula, with

$$D_i \mathcal{P}^i \equiv \frac{1}{2} D_i (\tilde{\psi} \gamma_h P^{ih} \psi) \equiv \frac{1}{2} D_i (\tilde{\psi} \gamma_h \psi) P^{ih} + \frac{1}{2} \tilde{\psi} \gamma_h \psi D_i P^{ih}. \quad (5.3)$$

The momentum constraint 0.2 gives

$$D_i P^{ih} = -T_0^h.$$

On the other hand, the identity 4.4 together with the Hamiltonian constraint implies that

$$D_i \mathcal{U}^i \equiv |D\psi|^2 - |\mathcal{D}\psi|^2 + \left(\frac{1}{2} T_{00} + \frac{1}{4} |K|^2 - \frac{1}{4} |\text{tr} K|^2\right) |\psi|^2. \quad (5.4)$$

We introduce the notations⁶

$$\nabla_i \psi := D_i \psi + \frac{1}{2} \gamma^h K_{ih} \psi, \quad \mathcal{N} \psi := \gamma^i \nabla_i \equiv (\mathcal{D} + \frac{1}{2} \text{tr} K) \psi. \quad (5.5)$$

Elementary computation using $D_i \eta^{hj} \equiv 0$, $D_i \gamma^h \equiv 0$ gives

$$|\nabla \psi|^2 := \eta^{ij} \widetilde{\nabla_i \psi} \nabla_j \psi \equiv |D\psi|^2 + \frac{1}{2} D_i (\tilde{\psi} \gamma_h \psi) K^{ih} + \frac{1}{4} |K|^2 |\psi|^2 \quad (5.6)$$

The identity 5.4 can therefore be written after simplification

$$D_i \mathcal{U}^i \equiv |\nabla \psi|^2 - |\mathcal{D}\psi|^2 + \left(\frac{1}{2} T_{00} - \frac{1}{4} |\text{tr} K|^2\right) |\psi|^2 - \frac{1}{2} D_i (\tilde{\psi} \gamma_h \psi) K^{ih}. \quad (5.7)$$

We deduce from the definition

$$|\mathcal{N} \psi|^2 \equiv |\mathcal{D}\psi|^2 + \frac{1}{2} D_i (\tilde{\psi} \gamma^i \psi) \text{tr} K + \frac{1}{4} |\text{tr} K|^2 |\psi|^2$$

which gives

$$D_i \mathcal{U}^i \equiv |\nabla \psi|^2 - |\mathcal{N} \psi|^2 + \frac{1}{2} T_{00} |\psi|^2 - \frac{1}{2} D_i (\tilde{\psi} \gamma^h \psi) P_{ih}. \quad (5.8)$$

Lemma 5 *If (M^n, g, K) is A.E. then it holds that*

$$D_i (\mathcal{U}^i + \mathcal{P}^i) \geq 0 \quad (5.9)$$

if the dominant energy condition holds and ψ satisfies the equation $\mathcal{N} \psi = 0$.

⁶Note that $\mathcal{N}$ is a linear operator mapping space spinor into space spinors, not the trace on M^n of the covariant derivative of a spacetime spinor.

Proof. The identities 5.3 and 5.8 lead to

$$D_i(\mathcal{U}^i + \mathcal{P}^i) \equiv |\nabla\psi|^2 - |\mathcal{N}\psi|^2 + \mathcal{T}, \quad \mathcal{T} := \frac{1}{2}(T_{00}|\psi|^2 - \tilde{\psi}\gamma^h\psi T_{0h}). \quad (5.10)$$

with $\mathcal{T} \geq 0$ under the dominant energy condition, because

$$|\tilde{\psi}\gamma^h\psi T_{0h}| \equiv |\psi|^2(\eta^{ih}T_{0i}T_{0h})^{\frac{1}{2}} \leq T_{00}|\psi|^2$$

with $\mathcal{T} \geq 0$ under the dominant energy condition, because

$$|\tilde{\psi}\gamma^h\psi T_{0h}| \equiv |\psi|^2(\eta^{ih}T_{0i}T_{0h})^{\frac{1}{2}} \leq T_{00}|\psi|^2.$$

■

Lemma 6 *If (M^n, g, K) is A.E., then the equation $\mathcal{N}\psi = 0$ has a solution $\psi \equiv \psi_0 + \psi_1$, ψ_0 smooth, constant in Ω_I and zero in the other ends, and $\psi_1 \in H_{s,\delta}$.*

Proof. The operator $\mathcal{N}$ has the same principal part as $\mathcal{D}$, therefore is also elliptic. It maps $H_{s,\delta}$ into $H_{s-1,\delta+1}$. The equation $\mathcal{N}\psi_1 = -\mathcal{N}\psi_0 \in H_{s-1,\delta+1}$ has one and only one solution if $\mathcal{N}$ is injective on $H_{s,\delta}$. To show injectivity⁷ we remark that the identity 5.10 was established without restriction on ψ , starting from the definitions of $\mathcal{U}^i$ and $\mathcal{P}^i$. We make $\psi = \psi_1$ in 5.10 and integrate it on M^n , the fall off of ψ_1 implies that the divergence gives no contribution, the equation $|\mathcal{N}\psi_1|^2 = 0$ implies therefore that on M^n , if $\mathcal{T} \geq 0$

$$|\nabla\psi_1|^2 = 0, \quad \text{i.e.} \quad D_i\psi_1 + \frac{1}{2}\gamma^h K_{ih}\psi_1 = 0, \quad \text{with} \quad \gamma^h K_{ih} \in H_{s-1,\delta+1}$$

The Poincaré inequality⁸ in weighted Hilbert spaces leads to $\psi_1 = 0$ if $\psi_1 \in H_{s,\delta}$, $s > \frac{n}{2} + 1$ and $-2 + \frac{n}{2} > \delta > -\frac{n}{2}$. ■

The lemmas, after choice of ψ_0 satisfying 5.2, prove the following theorem

Theorem 7 *If an Einsteinian spacetime satisfies the dominant energy condition, the energy momentum vector $E^0 = m$, $E^i = p^i$ of each end of an A.E. slice (M^n, g, K) satisfies the inequality*

$$m \geq |p|$$

⁷See a similar proof in Chrusciel Krakow lecture notes.

⁸See for instance Y. Choquet-Bruhat "General Relativity and the Einstein equations" appendix 3sobolev spaces" p.541

Acknowledgement 8 *I thank Thibault Damour for his encouragement to write this Note and for his help in doing it. I thank Piotr Chrusciel for communicating to me a text of his 2010 lectures in Krakow. A long list of references can be found there.*

16 Avenue d'Alembert, 92160, Antony
01 47 02 50 86, ycb@ihes.fr
Rubrique; Physique Mathématique